arXiv:1207.5891v2 [math.RA] 26 Jul 2012

DERIVATIONS AND IDENTITIES FOR FIBONACCI AND LUCAS POLYNOMIALS

LEONID BEDRATYUK

Анотація. We introduce the notion of Fibonacci and Lucas derivations of the polynomial algebras and prove that any element of kernel of the derivations defines a polynomial identity for the Fibonacci and Lucas polynomials. Also, we prove that any polynomial identity for Appel polynomial yields a polynomial identity for the Fibonacci and Lucas polynomials та описуємо arised interwining maps.

1. Introduction

The Fibonacci $F_n(x)$ and Lucas $L_n(x)$ polynomials are defined by the following ordinary generating functions

$$\mathcal{G}(F_n(x),t)=\frac{t}{1-xt-t^2}=\sum_{n=0}^{\infty}F_n(x)t^n,$$

$$\mathcal{G}(L_n(x),t)=\frac{1+t^2}{1-xt-t^2}=\sum_{n=0}^{\infty}L_n(x)t^n.$$

The derivatives of the polynomials have the form , see [1],[2]:

$$\frac{d}{dx}F_n(x)=\sum_{k=0}^{\left[\frac{n-1}{2}\right]}(-1)^k(n-1-2k)F_{n-1-2k}(x), \tag{1.1}$$

$$\frac{d}{dx}L_n(x)=n\sum_{k=0}^{\left[\frac{n-1}{2}\right]}(-1)^kL_{n-1-2k}(x), \tag{1.2}$$

We are interested in finding polynomial identities for the polynomials, i.e. identities of the form

$$P(F_0(x),F_1(x),\ldots,F_n(x))=\text{const} \;\text{ and } P(L_0(x),L_1(x),\ldots,L_n(x))=\text{const},$$

where $P(x_0,x_1,\ldots,x_n)$ is a polynomial of $n+1$ variables. We offer a method for finding such identities which is based on the followimng simple observation: if

$$\frac{d}{dx}P(F_0(x),F_1(x),\ldots,F_n(x))=0,$$

тоді $P(F_0(x),F_1(x),\ldots,F_n(x))=\text{const}$ i.e., it is an indentity.

On the other hand, rewrite this derivative in the form

$$\frac{d}{dx}P(F_0(x),F_1(x),\ldots,F_n(x))=$$

$$=\frac{\partial}{\partial x_0}P(x_0,x_1,\ldots,x_n)\Big|_{\{x_i=F_i(x)\}}\frac{d}{dx}F_0(x)+\cdots+\frac{\partial}{\partial x_n}P(x_0,x_1,\ldots,x_n)\Big|_{\{x_i=F_i(x)\}}\frac{d}{dx}F_n(x).$$

Therefore

$$\begin{gathered}\frac{d}{dx}P(F_0(x),F_1(x),\dots,F_n(x))=\\=\left(\frac{\partial}{\partial x_0}P(x_0,x_1,\dots,x_n)\mathcal{D}_{\mathcal{F}}(x_0)+\cdots+\frac{\partial}{\partial x_n}P(x_0,x_1,\dots,x_n)\mathcal{D}_{\mathcal{F}}(x_n)\right)\Big|_{\{x_i=F_i(x)\}}=\\=\mathcal{D}_{\mathcal{F}}(P(x_0,x_1,\dots,x_n))\Big|_{\{x_i=F_i(x)\}},\end{gathered}$$

where the differential operator $\mathcal{D}_{\mathcal{F}}$ defined by

$$\mathcal{D}_{\mathcal{F}}:=x_1\frac{\partial}{\partial x_2}+2x_2\frac{\partial}{\partial x_3}+(3\,x_3-x_1)\frac{\partial}{\partial x_4}+\cdots+\Big(\sum_{k=0}^{\left[\frac{n-1}{2}\right]}(-1)^k(n-1-2k)x_{n-1-2k})\Big)\frac{\partial}{\partial x_n}.$$

It is clear that if $\mathcal{D}_{\mathcal{F}}(P(x_0,x_1,\dots,x_n))$ then $\frac{d}{dx}P(F_0(x),F_1(x),\dots,F_n(x))=0$. Thus, any non-trivial polynomial $P(x_0,x_1,\dots,x_n)$, which belongs to the kernel of $\mathcal{D}_{\mathcal{F}}$, i.e., the following holds $\mathcal{D}_{\mathcal{F}}(P(x_0,x_1,\dots,x_n))=0$, defines a polynomial identity of the form $P(F_0(x),F_1(x),\dots,F_n(x))=$ const. Analogously, for the Lucas folynomials, we introduce the differential operator

$$\mathcal{D}_{\mathcal{L}}(x_n)=n\sum_{k=0}^{\left[\frac{n-1}{2}\right]}(-1)^k x_{n-1-2k}$$

and show that the condition

$$\mathcal{D}_{\mathcal{L}}(P(x_0,x_1,\dots,x_n))=0$$

implies the identity $P(L_0(x),L_1(x),\dots,L_n(x))=\text{const}$. For instance, it is easily verified that for the polynomial $x_1x_3-x_2^2$ we have $\mathcal{D}_{\mathcal{F}}(x_1x_3-x_2^2)=0$, thus $F_1(x)F_3(x)-F_2(x)^2$ is a constant. The substitution $F_n(x)$ of x_n gives

$$F_1(x)F_3(x)-F_2(x)^2=1.$$

A similar problem was solved in [3] for the Appel polynomials. In this paper was proved that any non-trivial element of kernel of the differential operator

$$\mathcal{D}_{\mathcal{A}}=x_0\frac{\partial}{\partial x_1}+2x_1\frac{\partial}{\partial x_2}+\cdots+nx_{n-1}\frac{\partial}{\partial x_n}.$$

defines some polynomial identity for the Appel polynomial. Recall that polynomials $\{A_n(x)\}$, $\deg(A_n(x))=n$ is called the Appel polynomials if

$$A_n'(x)=nA_{n-1}(x),n=0,1,2,\dots. \tag{1.3}$$

In the present paper we show how the known polylomial identities for Appel can be used to find a polynomials identities for the Fibonacci and Lucas polynomials. A linear map ψ_{AF} is called $(\mathcal{D}_{\mathcal{A}},\mathcal{D}_{\mathcal{F}})$-interwining map if the following condition holds: $\psi_{AF}\mathcal{D}_{\mathcal{A}}=\mathcal{D}_{\mathcal{F}}\psi_{AF}$. Any such map induces an ispmorphism from $\ker\mathcal{D}_{\mathcal{A}}$ to $\ker\mathcal{D}_{\mathcal{F}}$. For instance, the discriminant of the polynomial (in the variables X,Y)

$$x_0X^3+3\,x_1X^2Y+3\,x_2XY^2+x_3Y^3,$$

equals

$$\begin{vmatrix} x_0 & 3x_1 & 3x_2 & x_3 & 0 \\ 0 & x_0 & 3x_1 & 3x_2 & x_3 \\ 3x_0 & 6x_1 & 3x_2 & 0 & 0 \\ 0 & 3x_0 & 6x_1 & 3x_2 & 0 \\ 0 & 0 & 3x_0 & 6x_1 & 3x_2 \end{vmatrix} = 27(6\,x_0x_3x_2x_1 + 3\,{x_1}^2{x_2}^2 - 4\,{x_1}^3x_3 - 4\,{x_2}^3x_0 - {x_0}^2{x_3}^2),$$

and lies in the kernel of the operator $\mathcal{D}_{\mathcal{A}}$. It is well known result of the classical invariant theory. It is easily checked that the linear map defined by

$$\psi_{AL}(x_0) = x_0, \psi_{AL}(x_1) = x_1,$$
$$\psi_{AL}(x_2) = x_2, \psi_{AL}(x_3) = x_3 + 3x_1,$$

commutes with the operators $\mathcal{D}_{\mathcal{A}}$ and $\mathcal{D}_{\mathcal{L}}$. Therefore the element

$$\begin{vmatrix} \psi_{AL}(x_0) & 3\psi_{AL}(x_1) & 3\psi_{AL}(x_2) & \psi_{AL}(x_3) & 0 \\ 0 & \psi_{AL}(x_0) & 3\psi_{AL}(x_1) & 3\psi_{AL}(x_2) & \psi_{AL}(x_3) \\ 3\psi_{AL}(x_0) & 6\psi_{AL}(x_1) & 3\psi_{AL}(x_2) & 0 & 0 \\ 0 & 3\psi_{AL}(x_0) & 6\psi_{AL}(x_1) & 3\psi_{AL}(x_2) & 0 \\ 0 & 0 & 3\psi_{AL}(x_0) & 6\psi_{AL}(x_1) & 3\psi_{AL}(x_2) \end{vmatrix} = \begin{vmatrix} x_0 & 3x_1 & 3x_2 & x_3+3x_1 & 0 \\ 0 & x_0 & 3x_1 & 3x_2 & x_3+3x_1 \\ 3x_0 & 6x_1 & 3x_2 & 0 & 0 \\ 0 & 3x_0 & 6x_1 & 3x_2 & 0 \\ 0 & 0 & 3x_0 & 6x_1 & 3x_2 \end{vmatrix},$$

lies in the kernel of the operator $\mathcal{D}_{\mathcal{L}}$ and defines the following identity for the Lucas polynomial:

$$\begin{vmatrix} L_0(x) & 3L_1(x) & 3L_2(x) & L_3(x)+3L_1(x) & 0 \\ 0 & L_0(x) & 3L_1(x) & 3L_2(x) & L_3(x)+3L_1(x) \\ 3L_0(x) & 6L_1(x) & 3L_2(x) & 0 & 0 \\ 0 & 3L_0(x) & 6L_1(x) & 3L_2(x) & 0 \\ 0 & 0 & 3L_0(x) & 6L_1(x) & 3L_2(x) \end{vmatrix} = -864.$$

In the paper we present methods of the theory of locally nilpotent derivation to find polynomial identities for the Fibonacci and Lucal polynomials.

In section 2,we give a brief introduction to the theory of localy nilpotent derivations. Also, we introduce the notion of the Fibonacci and Lucas derivations and find its kernels. In this way we obtain some polynomials identities for the Fibonacci and Lucas polynomials.

In the section 3 we find a $(\mathcal{D}_{\mathcal{A}}, \mathcal{D}_{\mathcal{F}})$-interwining map and a $(\mathcal{D}_{\mathcal{A}}, \mathcal{D}_{\mathcal{L}})$-interwining map.

2. Fibonacci and Lucas derivations

2.1. **Derivations and its kernels.** Let $\mathbb{C}[x_0, x_1, x_2, \ldots, x_n]$ be the polynomial algebra in $n+1$ variables $x_0, x_1, x_2, \ldots, x_n$ over $\mathbb{C}$. Recall that a *derivation* of the polynomial algebra $\mathbb{C}[x_0, x_1, x_2, \ldots, x_n]$ is a linear map D satisfying the Leibniz rule:

$$D(x_1\,x_2) = D(x_1)x_2 + x_1D(x_2), \text{ for all } x_1, x_2 \in \mathbb{C}[x_0, x_1, x_2, \ldots, x_n].$$

A derivation D is called *locally nilpotent* if for every $f \in \mathbb{C}[x_0, x_1, x_2, \ldots, x_n]$ there is an $n \in \mathbb{N}$ such that $D^n(f) = 0$. Any derivation D is completely determined by the elements $D(x_i)$. A derivation D is called *linear* if $D(x_i)$ is a linear form. A linear locally nilpotent derivation is called a *Weitzenböck derivation.* A derivation D is called the triangular if $D(x_i) \in \mathbb{C}[x_0, \ldots, x_{i-1}]$. Any triangular derivation is locally nilpotent.

The subalgebra

$$\ker D := \{f \in \mathbb{C}[x_0, x_1, x_2, \ldots, x_n] \mid D(f) = 0\},$$

is called the *kernel* of derivation D.

For arbitrary localy niplotent derivation D tye following statement holds:

Theorem 2.1. *Suppose that there exists a polynomials h such that $D(h) \neq 0$ but $D^2(h) = 0$. Then*

$$\ker D = \mathbb{C}[\sigma(x_0), \sigma(x_2), \ldots, \sigma(x_n)][D(h)^{-1}] \cap \mathbb{C}[x_0, x_1, \ldots, x_n],$$

where σ is the Diximier map

$$\sigma(x_i) = \sum_{k=0}^{\infty} D^k(x_i) \frac{\lambda^k}{k!}, \lambda = -\frac{h}{D(h)}, D(\lambda) = -1.$$

The proof one may find in [5] and [6].

The expressions (1.1),(1.2) motivate the following definition

Definition 1. *Derivations of $\mathbb{C}[x_0, x_1, x_2, \ldots, x_n]$ defined by*

$$D_{\mathcal{F}}(x_0) = D_{\mathcal{F}}(x_1) = 0, D_{\mathcal{F}}(x_n) = \sum_{k=0}^{\left[\frac{n-1}{2}\right]} (-1)^k (n-1-2k) x_{n-1-2k},$$

$$D_{\mathcal{L}}(x_0) = 0, D_{\mathcal{L}}(x_n) = n \sum_{k=0}^{\left[\frac{n-1}{2}\right]} (-1)^k x_{n-1-2k}, i = 2, 3, \ldots, n, \ldots,$$

are called **the Fibonacci derivation** *and* **the Lucas derivation** *respectivelly*

We have

$$\begin{array}{ll}
D_{\mathcal{F}}(x_0) = 0, & D_{\mathcal{L}}(x_0) = 0,\\
D_{\mathcal{F}}(x_1) = 0, & D_{\mathcal{L}}(x_1) = x_0,\\
D_{\mathcal{F}}(x_2) = x_1, & D_{\mathcal{L}}(x_2) = 2x_1\\
D_{\mathcal{F}}(x_3) = 2x_2, & D_{\mathcal{L}}(x_3) = 3\,x_2 - 3\,x_0,\\
D_{\mathcal{F}}(x_4) = 3\,x_3 - x_1, & D_{\mathcal{L}}(x_4) = 4\,x_3 - 4\,x_1,\\
D_{\mathcal{F}}(x_5) = 4\,x_4 - 2\,x_2, & D_{\mathcal{L}}(x_5) = 5\,x_4 - 5\,x_2\\
D_{\mathcal{F}}(x_6) = 5\,x_5 - 3\,x_3 + x_1, & D_{\mathcal{L}}(x_6) = 6\,x_5 - 6\,x_3 + 6x_1.
\end{array}$$

We define the substitution homomorphisms $\varphi_{\mathcal{F}}, \varphi_{\mathcal{L}} : \mathbb{C}[x_0, x_1, \ldots, x_n] \to \mathbb{C}[x]$ by $\varphi_{\mathcal{F}}(x_i) = F_i(x)$ and by $\varphi_{\mathcal{L}}(x_i) = L_i(x)$. Put

$$\ker \varphi_{\mathcal{F}} := \{P(x_0, x_1, ..., x_n) \mid \varphi_{\mathcal{F}}(P(x_0, x_1, ..., x_n)) = 0\},$$
$$\ker \varphi_{\mathcal{L}} := \{P(x_0, x_1, ..., x_n) \mid \varphi_{\mathcal{L}}\,(P(x_0, x_1, ..., x_n)) = 0\}.$$

Any element of $\ker \varphi_{\mathcal{F}}$ or $\ker \varphi_{\mathcal{F}}$ defines a polynomial identity for the Fibonacci and Lucas polynomials відповідно. Put

$$\ker \mathcal{D}_{\mathcal{F}} := \{S \in \mathbb{C}[x_0, x_1, ..., x_n] \mid \mathcal{D}_{\mathcal{F}}(S) = 0\},$$
$$\ker \mathcal{D}_{\mathcal{L}} := \{S \in \mathbb{C}[x_0, x_1, ..., x_n] \mid \mathcal{D}_{\mathcal{L}}(S) = 0\}.$$

It is easy to see that $\varphi_{\mathcal{F}} \mathcal{D}_{\mathcal{F}} = \frac{d}{dx} \varphi_{\mathcal{F}}$ and $\varphi_{\mathcal{L}} \mathcal{D}_{\mathcal{L}} = \frac{d}{dx} \varphi_{\mathcal{L}}$. It follows that $\varphi_{\mathcal{F}}(\ker \mathcal{D}_{\mathcal{F}}) \subset \ker \varphi_{\mathcal{F}}$ and $\varphi_{\mathcal{L}}(\ker \mathcal{D}_{\mathcal{L}}) \subset \ker \varphi_{\mathcal{L}}$. Note that $\varphi_{\mathcal{L}}(\ker \mathcal{D}_{\mathcal{L}}) \neq \ker \varphi_{\mathcal{L}}$. In fact, we have $\varphi_{\mathcal{F}}(x_n - x_2 x_{n-1} + x_{n-2}) = F_n(x) - (F_2(x) F_{n-1}(x) + F_{n-2}(x)) = 0$ but $x_n - x_2 x_{n-1} - x_{n-2} \notin \ker \mathcal{D}_{\mathcal{F}}$.

We have thus proved the following theorem.

Theorem 2.2. *Let $P(x_0, x_1, \ldots, x_n)$ be a polynomial.*

(i) If $\mathcal{D}_{\mathcal{F}}(P(x_0, x_1, \ldots, x_n)) = 0$ then $P(F_0(x), F_1(x), \ldots, F_n(x)) = \mathrm{const}$;

(ii) if $\mathcal{D}_{\mathcal{L}}(P(x_0, x_1, \ldots, x_n)) = 0$ then $P(L_0(x), L_1(x), \ldots, L_n(x)) = \mathrm{const}$.

2.2. **The kernel of the Fibonacci derivation.** It is obviously that this derivation is triangular and hence localy nilpotent. Thus to find its kernel we may use Theorem 2.1.

Let us construct the Diximier map for the Fibonacci derivation. For this purpose, we derive first a close expression for $D_{\mathcal{F}}^k(x_n)$.

We have

$$D_{\mathcal{F}}^2(x_n) = D_{\mathcal{F}}\left(\sum_{k=0}^{\left[\frac{n-1}{2}\right]}(-1)^k(n-1-2k)x_{n-1-2k}\right) =$$

$$= \sum_{k=0}^{\left[\frac{n-1}{2}\right]}(-1)^k(n-1-2k)\sum_{j=0}^{\left[\frac{n-2-2k}{2}\right]}(-1)^j(n-2-2k-2j)x_{n-2-2j} =$$

$$= \sum_{i=0}^{\left[\frac{n-2}{2}\right]}(-1)^i(i+1)(n-2-2i)(n-(i+1))x_{n-2-2i}.$$

Similarly we get

$$D_{\mathcal{F}}^3(x_n) = \sum_{i=0}^{\left[\frac{n-3}{2}\right]}(-1)^i\frac{(i+1)(i+2)}{2}(n-3-2i)(n-(i+1))(n-(i+2))x_{n-3-2i}.$$

Induction gives

$$D_{\mathcal{F}}^k(x_n) = (k-1)!\sum_{i=0}^{\left[\frac{n-k}{2}\right]}(-1)^i(n-k-2i)\binom{i+k-1}{k-1}\binom{n-i-1}{k-1}x_{n-k-2i}.$$

Since $D_{\mathcal{F}}\left(-\frac{x_2}{x_1}\right) = -1$ we put $\lambda = -\frac{x_2}{x_1}$. Now we may find the Diximier map:

$$\sigma(x_n) = \sum_{k=0}^{n-1}D_{\mathcal{F}}^k(x_n)\frac{\lambda^k}{k!} =$$

$$= x_n + \sum_{k=1}^{n-1}\frac{\lambda^k}{k}\sum_{i=0}^{\left[\frac{n-k}{2}\right]}(-1)^{k+i}(n-k-2i)\binom{i+k-1}{k-1}\binom{n-i-1}{k-1}x_{n-k-2i} =$$

$$= x_n + \sum_{k=1}^{n-3}\frac{\lambda^k}{k}\sum_{i=0}^{\left[\frac{n-k}{2}\right]}(-1)^i(n-k-2i)\binom{i+k-1}{k-1}\binom{n-i-1}{k-1}x_{n-k-2i} + (n-1)x_2\lambda^{n-2} + x_1\lambda^{n-1}.$$

Replasing λ by $-\frac{x_2}{x_1}$, we obtain, after simplifying:

$$x_1^{n-2}\sigma(x_n) =$$

$$= x_nx_1^{n-2} + \sum_{k=1}^{n-3}\frac{1}{k}\sum_{i=0}^{\left[\frac{n-k}{2}\right]}(-1)^{k+i}(n-k-2i)\binom{i+k-1}{k-1}\binom{n-i-1}{k-1}x_{n-k-2i}x_2^kx_1^{n-2-k} + (n-2)(-1)^{n-2}x_2^{n-1}.$$

The polynomials

$$
\begin{aligned}
C_n :=& \\
&= x_n x_1^{n-2} + \sum_{k=1}^{n-3} \frac{1}{k} \sum_{i=0}^{\left[\frac{n-k}{2}\right]} (-1)^{k+i} (n-k-2i) \binom{i+k-1}{k-1} \binom{n-i-1}{k-1} x_{n-k-2i} x_2^k x_1^{n-2-k} + \\
&\qquad + (n-2)(-1)^{n-2} x_2^{n-1}, n > 2,
\end{aligned}
$$

belong to the kernel $\ker \mathcal{D}_{\mathcal{F}}$. We call them *the Cayley elements* of the localy nilpotent derivation $\mathcal{D}_{\mathcal{F}}$. The first few Cayley polynomials are shown below:

$$
\begin{aligned}
C_3 &= -{x_2}^2 + x_3 x_1, \\
C_4 &= 2\,{x_2}^3 - 3\,x_2 x_3 x_1 + {x_1}^2 x_2 + x_4 {x_1}^2, \\
C_5 &= -3\,{x_2}^4 + 6\,{x_2}^2 x_3 x_1 - {x_1}^2 {x_2}^2 - 4\,x_2 x_4 {x_1}^2 + x_5 {x_1}^3, \\
C_6 &= 4\,{x_2}^5 - 10\,{x_2}^3 x_3 x_1 - 2\,{x_1}^2 {x_2}^3 + 10\,{x_2}^2 x_4 {x_1}^2 - 5\,x_2 x_5 {x_1}^3 + 3\,{x_1}^3 x_2 x_3 - {x_1}^4 x_2 + x_6 {x_1}^4.
\end{aligned}
$$

Theorem 2.1 implies

Theorem 2.3.

$$\ker \mathcal{D}_{\mathcal{F}} = \mathbb{C}[x_0, x_1, C_3, C_4, \ldots, C_n][x_1^{-1}] \cap \mathbb{C}[x_0, x_1, \ldots, x_n].$$

Thus, we obtain a description of the kernel of the Fibonacci derivation.

To get an identity for the Fibonacci polynomials we should find $\varphi_F(C_n)$. We get

$$
\begin{aligned}
\varphi_F(C_n) &= \varphi_F(C_n(x_0, x_1, \ldots, x_n)) = F_n(x) + \\
&+ \sum_{k=1}^{n-3} \frac{1}{k} \sum_{i=0}^{\left[\frac{n-k}{2}\right]} (-1)^{k+i} (n-k-2i) \binom{i+k-1}{k-1} \binom{n-i-1}{k-1} F_{n-k-2i}(x) F_2(x)^k + \\
&\qquad + (n-2)(-1)^{n-2} F_2(x)^{n-1} = \text{const}.
\end{aligned}
$$

Conjecture. $\varphi_F(C_n) = C_n(F_1(x), \ldots, F_n(x)) = \begin{cases} 0, & n \text{ even}, \\ 1, & n \text{ odd}. \end{cases}$

2.3. **Ядро диференціювання Люка.** Для диференціювання $D_{\mathcal{L}}$ отримаємо схожим способом

$$D_{\mathcal{L}}^k(x_n) = n\,(k-1)! \sum_{i=0}^{\left[\frac{n-k}{2}\right]} (-1)^i \binom{i+k-1}{k-1} \binom{n-i-1}{k-1} x_{n-k-2i}.$$

Маємо, що $D_{\mathcal{L}}\left(-\dfrac{x_1}{x_0}\right) = -1$. Шукаємо відображення Діксм'є:

$$\sigma(x_n) = \sum_{k=0}^{n} D_{\mathcal{L}}(x_n) \frac{\lambda^i}{k!} =$$

$$= x_n + n \sum_{k=1}^{n-2} \frac{\lambda^k}{k} \sum_{n-k-2i>0} (-1)^{k+i} \binom{i+k-1}{k-1} \binom{n-i-1}{k-1} x_{n-k-2i} + n x_1 \lambda^{n-1} + x_0 \lambda^n$$

Відповідний елемент Келлі має вигляд

$$\sigma_n(x_n) = x_n x_0^{n-1} + n\sum_{k=1}^{n-2}\frac{1}{k}\sum_{i=0}^{\left[\frac{n-k}{2}\right]}(-1)^{k+i}\binom{i+k-1}{k-1}\binom{n-i-1}{k-1}x_{n-k-2i}x_1^k x_0^{n-1-k} + (n-1)(-1)^{n-1}x_1^n,$$

Theorem 2.4.

$$k(x_0, x_1, \ldots, x_n)^{\mathcal{D}_{\mathcal{L}}} = k[x_0, C_2, C_3, C_4, \ldots, C_n][x_0^{-1}],$$
$$k[x_0, x_1, \ldots, x_n]^{\mathcal{D}_{\mathcal{L}}} = k[x_0, C_2, C_3, C_4, \ldots, C_n][x_0^{-1}] \cap k[x_0, x_1, \ldots, x_n].$$

Випишемо після спрощення декілька цих многочленів

$$C_1 = x_0,$$
$$C_2 = x_2x_0 - {x_1}^2,$$
$$C_3 = 2\,{x_1}^3 + 3\,x_1{x_0}^2 - 3\,x_1x_2x_0 + x_3{x_0}^2,$$
$$C_4 = -3\,{x_1}^4 - 4\,{x_1}^2{x_0}^2 + 6\,{x_1}^2x_2x_0 - 4\,x_1x_3{x_0}^2 + x_4{x_0}^3,$$
$$C_5 = 4\,{x_1}^5 + 10\,{x_1}^2x_3{x_0}^2 - 5\,x_1{x_0}^4 - 5\,x_1x_4{x_0}^3 - 10\,{x_1}^3x_2x_0 + 5\,x_1x_2{x_0}^3 + x_5{x_0}^4.$$

Гіпотеза. $C_n(L_0(x), \ldots, L_n(x)) = \begin{cases} 2, & n - \text{ парне} \\ 0, & \text{інакше} \end{cases}$

3. Appel-Lucas and Appel-Fibonacci interwining maps

3.1. **Appel-Lucas interwinning map.** Denote by ψ_{AL} a Appel-Lucas intewinning map. Будемо шукати його у вигляді:

$$\psi_{AL}(x_n) = x_n + \alpha_n^{(1)}x_{n-2} + \alpha_n^{(2)}x_{n-4} + \ldots + \alpha_n^{(i)}x_{n-2i} + \ldots + \alpha_n^{(\left[\frac{n-1}{2}\right])}x_{n-2\left[\frac{n-1}{2}\right]}.$$

Доведемо таке твердження

Lemma 3.1. *Послідовності* $\alpha_n^{(1)}$, $\alpha_n^{(2)}, \ldots, \alpha_n^{(\left[\frac{n-1}{2}\right])}$ *задовільняють такій системи рекурентних рівнянь*

$$\begin{cases} (n-2)\alpha_n^{(1)} = n(\alpha_{n-1}^{(1)} + 1), \\ (n-4)\alpha_n^{(2)} = n(\alpha_{n-1}^{(2)} + \alpha_{n-1}^{(1)}), \\ (n-2i)\alpha_n^{(i)} = n(\alpha_{n-1}^{(i)} + \alpha_{n-1}^{(i-1)}), \\ \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots \\ (n-2\left[\frac{n-1}{2}\right])\alpha_n^{(\left[\frac{n-1}{2}\right])} = n\left(\alpha_{n-1}^{(\left[\frac{n-1}{2}\right])} + \alpha_{n-1}^{(\left[\frac{n-1}{2}\right]-1)}\right), \end{cases}$$

із початковими умовами $\alpha_{2i}^{(i)} = 0$.

Доведення. We have

$$D_{\mathcal{L}}(\psi_{AL}(x_n)) = n(x_{n-1} - x_{n-3} + x_{n-5} - x_{n-7} + \ldots) + \alpha_n^{(1)}(n-2)(x_{n-3} - x_{n-5} + \ldots) +$$
$$+ \alpha_n^{(2)}(n-4)(x_{n-5} - x_{n-7} + \ldots) + \ldots$$

Звідси знаходимо

$$D_{\mathcal{L}}(\psi_{AL}(x_n)) = nx_{n-1} + x_{n-3}((n-2)\alpha_n^{(1)} - n) + x_{n-5}((n-4)\alpha_n^{(2)} - (n-2)\alpha_n^{(1)} + n) +$$
$$+ x_{n-2i-1}((n-2i)\alpha_n^{(i)} - (n-2(i-1))\alpha_n^{(i-1)} + (n-2(i-2))\alpha_n^{(i-2)} + \ldots) + \ldots +$$

З іншого боку, має виконуватися тотожність,

$$D_{\mathcal{L}}(\psi_{AL}(x_n)) = \psi_{AL}(\mathcal{D}(x_n)) = n\psi_{AL}(x_{n-1}) = n \sum_{n-1-2i>0} \alpha_{n-1}^{(i)} x_{n-1-2i}(-1)^i.$$

Прирівнюючи відповідні коефіцієнти, отримуємо такі рекурентні співвідношення на $\alpha_n^{(i)}$:

$$(n-2)\alpha_n^{(1)} - n = n\alpha_{n-1}^{(1)}, \alpha_2^{(1)} = 0,$$
$$(n-4)\alpha_n^{(2)} - (n-2)\alpha_n^{(1)} + n = n\alpha_{n-1}^{(2)}, \alpha_4^{(1)} = 0,$$
$$\dots$$
$$\sum_{n-2i>0} (n-2i)\alpha_n^{(i)}(-1)^i = n\alpha_{n-1}^{(i)}, \alpha_{2i}^{(i)} = 0.$$

Після спрощення отримуємо систему рекурентних рівнянь

$$\begin{cases} (n-2)\alpha_n^{(1)} = n(\alpha_{n-1}^{(1)} + 1), \\ (n-4)\alpha_n^{(2)} = n(\alpha_{n-1}^{(2)} + \alpha_{n-1}^{(1)}), \\ (n-2i)\alpha_n^{(i)} = n(\alpha_{n-1}^{(i)} + \alpha_{n-1}^{(i-1)}), \\ \dots\dots\dots\dots\dots\dots\dots\dots\dots\dots \\ (n-2\left[\frac{n-1}{2}\right])\alpha_n^{(\left[\frac{n-1}{2}\right])} = n\left(\alpha_{n-1}^{(\left[\frac{n-1}{2}\right])} + \alpha_{n-1}^{(\left[\frac{n-1}{2}\right]-1)}\right), \end{cases}$$

на послідовності $\alpha_n^{(1)}, \alpha_n^{(2)}, \dots, \alpha_n^{(\left[\frac{n-1}{2}\right])}$ із початковими умовами $\alpha_{2i}^{(i)} = 0$.

□

Розглянемо допоміжне неоднорідне рекурентне співвідношення

$$(n-a)x_n = n(x_{n-1} + g_{n-1}), x_a = 0, n \ge a.$$

а g_n – деяка фіксована послідовність. Тоді

Lemma 3.2.

$$x_n = n^{\underline{a}} \sum_{i=a}^{n-1} \frac{g_i}{i^{\underline{a}}} = \binom{n}{a} \sum_{i=a}^{n-1} \frac{g_i}{\binom{i}{a}}$$

де $n^{\underline{a}} := n(n-1)(n-2)\cdots(n-(a-1))$

Доведення. Справді, нехай

$$x_n = n^{\underline{a}} \sum_{i=a}^{n-1} \frac{g_i}{i^{\underline{a}}}.$$

Тоді

$$n(x_{n-1} + g_{n-1}) = n\left((n-1)^{\underline{a}} \sum_{i=a}^{n-2} \frac{g_i}{i^{\underline{a}}} + g_{n-1}\right) =$$
$$= n\left((n-1)^{\underline{a}} \sum_{i=a}^{n-2} \frac{g_i}{i^{\underline{a}}} + \frac{(n-1)^{\underline{a}} g_{n-1}}{(n-1)^{\underline{a}}}\right) = n(n-1)^{\underline{a}} \left(\sum_{i=a}^{n-2} \frac{g_i}{i^{\underline{a}}} + \frac{g_{n-1}}{(n-1)^{\underline{a}}}\right) =$$
$$= n(n-1)^{\underline{a}} \sum_{i=a}^{n-1} \frac{g_i}{i^{\underline{a}}} = (n-a)n^{\underline{a}} \sum_{i=a}^{n-1} \frac{g_i}{i^{\underline{a}}} = (n-a)x_n.$$

□

Як наслідок маємо, що система рекурентних рівнянь має такий розвязок

$$\alpha_n^{(1)} = n(n-2),$$
$$\alpha_n^{(k)} = (n)_{2k-1} \sum_{i=2k}^{n-1} \frac{\alpha_i^{(k-1)}}{(i)_{2k-1}}.$$

Розв'язуючи систему послідовно знаходимо

$$\begin{aligned}
\alpha_n^{(1)} &= n(n-2),\\
\alpha_n^{(2)} &= \frac{1}{4}\,(n-1)\,n\,(3\,n-7)\,(n-4) = \frac{1}{2!}(n-4)\binom{n}{2}(3n-7),\\
\alpha_n^{(3)} &= \frac{1}{3!}(n-6)\binom{n}{3}\left(19\,n^2-141\,n+254\right),\\
\alpha_n^{(4)} &= \frac{1}{4!}(n-8)\binom{n}{4}\left(211\,n^3-3258\,n^2+16481\,n-27306\right),\\
\alpha_n^{(5)} &= \frac{1}{5!}(n-10)\binom{n}{5}\left(3651\,n^4-96550\,n^3+946185\,n^2-4071950\,n+6492024\right)
\end{aligned}$$

Для знаходження загального розв'язку системи розглянемо невідомі послідовності $\alpha_n^{(s)}$ у базисі із спадних факторіалів. Нехай

$$\alpha_n^{(s)} = \beta_0^{(s)} n^{\underline{s}} + \beta_1^{(s)} n^{\underline{s+1}} + \cdots + \beta_s^{(s)} n^{\underline{2s}},$$

Легко бачити, що

$$(n-s)n^{\underline{s}} = n^{\overline{s+1}}, n(n-1)^{\underline{s}} = n^{\overline{s+1}}.$$

Тоді

$$(n-2s)\alpha_n^{(s)} = (n-2s)(\beta_0^{(s)} n^{\underline{s}} + \beta_1^{(s)} n^{\underline{s+1}} + \cdots + \beta_s^{(s)} n^{\underline{2s}}) =$$
$$= ((n-s)-s)\beta_0^{(s)} n^{\overline{s}} + ((n-(s+1))-(s-1)))\beta_1^{(s)} n^{\underline{s+1}} + \cdots + (n-2s)\beta_s^{(s)} n^{\underline{2s}} =$$
$$= \sum_{i=0}^{s} \beta_i^{(s)} n^{\overline{s+i+1}} - \sum_{i=0}^{s-1} (s-i)\beta_i^{(s)} n^{\overline{s+i}}$$

З іншого боку

$$n(\alpha_{n-1}^{(s)} + \alpha_{n-1}^{(s-1)}) = \sum_{i=0}^{s} n\beta_i^{(s)}(n-1)^{\overline{s+i}} + \sum_{i=0}^{s-1} n\beta_i^{(s-1)}(n-1)^{\overline{s-1+i}} = \sum_{i=0}^{s} \beta_i^{(s)} n^{\overline{s+i+1}} + \sum_{i=0}^{s-1} \beta_i^{(s-1)} n^{\overline{s+i}}.$$

Звідси, прирівнявши відповідні коефіцієнти, зразу знаходимо, що

$$\beta_i^{(s)} = \frac{\beta_i^{(s-1)}}{i-s}, i = 0 \ldots s-1.$$

Коефіцієнт $\beta_s^{(s)}$ знайдемо з умови, що $a_{2s}^{(s)} = 0$. Маємо

$$\sum_{i=0}^{s-1} \frac{\beta_i^{(s-1)}}{i-s}(2s)^{\overline{s+i}} + \beta_s^{(s)}(2s)! = 0$$

Звідси

$$\beta_s^{(s)} = \frac{1}{(2s)!} \sum_{i=0}^{s-1} \frac{\beta_i^{(s-1)}}{s-i} \overline{(2s)}^{s+i} = \sum_{i=0}^{s-1} \frac{\beta_i^{(s-1)}}{(s-i)(s-i)!} = -\sum_{i=0}^{s-1} \frac{\beta_i^{(s)}}{(s-i)!}.$$

Таким чином, маємо такі рекурентні співвідношення для $\beta_n^{(s)}$:

$$\begin{aligned}
\beta_0^{(s)} &= -\frac{\beta_0^{(s-1)}}{s}, \\
\beta_1^{(s)} &= -\frac{\beta_1^{(s-1)}}{s-1}, \\
\beta_2^{(s)} &= -\frac{\beta_1^{(s-1)}}{s-2}, \\
\dots \\
\beta_{s-1}^{(s)} &= -\beta_1^{(s-1)}, \\
\beta_s^{(s)} &:= b_s = -\sum_{i=0}^{s-1} \frac{\beta_i^{(s)}}{(s-i)!}.
\end{aligned}$$

Звідси знаходимо, що

$$\beta_i^{(s)} = \frac{(-1)^{s-i}}{(s-i)!} b_i, \text{ for } i = 0, \dots, s-1,$$

Отримуємо таке рекурентне співвідношення на послідовність b_s :

$$b_s = -\frac{(-1)^s}{(s!)^2} - \sum_{i=1}^{s-1} \frac{(-1)^{s-i}}{((s-i)!)^2} b_i,$$

Поклавши $b_0 = 1$ отримаємо, що елементи послідовності b_n задовільняють таке рекурентне співвідношення

$$\sum_{i=0}^{n} \frac{(-1)^{n-i}}{(n-i)!^2} b_n = 0, n > 0$$

Розглянемо ряд

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{n!^2} z^n = J_0(\sqrt{4z}\,)$$

і звичайну породжуючу функцію

$$G(b_n, z) = \sum_{n=0}^{\infty} b_n x^n.$$

Тоді, враховуючи рекурентне співвідношення знайдемо, що

$$J_0(\sqrt{4z}\,) G(b_n, z) = 1.$$

Звідси

$$G(b_n, z) = \frac{1}{J_0(\sqrt{4z}\,)} = 1 + z + \frac{3}{4} z^2 + \frac{19}{36} z^3 + \frac{211}{576} z^4 + \frac{1217}{4800} z^5 + \frac{30307}{172800} z^6 + \cdots$$

Отже ми довели таку теорему

Theorem 3.1. *A Appel-Lucas interwining map has the form*

$$\psi_{AL}(x_n) = x_n + \alpha_n^{(1)} x_{n-2} + \alpha_n^{(2)} x_{n-4} + \ldots + \alpha_n^{(i)} x_{n-2i} + \ldots + \alpha_n^{([\frac{n-1}{2}])} x_{n-2[\frac{n-1}{2}]},$$

where

$$\alpha_n^{(s)} = \frac{(-1)^s}{s!} b_0 n^{\underline{s}} + \cdots + \frac{(-1)^{s-i}}{(s-i)!}\, b_i n^{\underline{s+i}} + \cdots + b_s n^{\underline{2s}},$$

and the generating function for $b_0, b_1, \ldots, b_n, \ldots$ are defined через обернену функцію Бесселя

$$\sum_{i=0}^{\infty} b_i z^i = J_0^{-1}(\sqrt{4z}\,).$$

3.2. **Appel-Fibonacci interwining map.** Шукаємо явний вигляд ізоморфізму ядер $\ker \mathcal{D}_{\mathcal{A}} \overset{\psi}{\to} \ker \mathcal{D}_{\mathcal{F}}$.

Шукаємо ψ у вигляді:

$$\psi(x_n) = \alpha_n^{(0)} x_{n+1} + \alpha_n^{(1)} x_{n-1} + \alpha_n^{(2)} x_{n-3} + \ldots + \alpha_n^{(i)} x_{n+1-2i} + \ldots + \alpha_n^{([\frac{n-1}{2}])} x_{n+1-2[\frac{n-1}{2}]}.$$

Доведемо твердження

Lemma 3.3. *Коефіцієнти $\alpha_n^{(1)}, \alpha_n^{(2)}, \ldots, \alpha_n^{([\frac{n-1}{2}])}$ задовільняють такій системи рекурентних рівнянь*

$$\begin{cases} \alpha_n^{(0)} = 1, \\ \alpha_n^{(1)} = n\left(\dfrac{\alpha_{n-1}^{(1)}}{n-2} + \dfrac{\alpha_{n-1}^{(0)}}{n}\right), \\ \alpha_n^{(2)} = n\left(\dfrac{\alpha_{n-1}^{(2)}}{n-4} + \dfrac{\alpha_{n-1}^{(1)}}{n-2}\right), \\ \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots \\ \alpha_n^{(s)} = n\left(\dfrac{\alpha_{n-1}^{(s)}}{n-2s} + \dfrac{\alpha_{n-1}^{(s-1)}}{n-2(s-1)}\right), \\ \ldots \end{cases}$$

із початковими умовами $\alpha_{2i}^{(i)} = 0$.

Доведення. Тоді

$$D_{\mathcal{F}}(\psi(x_n)) = \alpha_n^{(0)}(nx_n - (n-2)x_{n-2} + (n-4)x_{n-4} - (n-6)x_{n-6} + \ldots) +$$
$$+ \alpha_n^{(1)}((n-2)x_{n-2} - (n-4)x_{n-4} + \ldots) + \alpha_n^{(2)}((n-4)x_{n-4} - (n-6)x_{n-6} + \ldots) + \ldots$$

Звідси знаходимо

$$D_{\mathcal{F}}(\varphi(x_n)) = n\alpha_n^{(0)} x_n + (n-2)x_{n-2}(\alpha_n^{(1)} - \alpha_n^{(0)}) + (n-4)x_{n-4}(\alpha_n^{(2)} - \alpha_n^{(1)} + \alpha_n^{(0)}) +$$
$$+ x_{n-2i}(n-2i)(\alpha_n^{(i)} - \alpha_n^{(i-1)} + \alpha_n^{(i-2)} + \cdots + (-1)^i \alpha_n^{(0)}) + \ldots +$$

З іншого боку, має виконуватися тотожність,

$$D_{\mathcal{F}}(\psi(x_n)) = \psi(\mathcal{D}(x_n)) = n\psi(x_{n-1}) = n\sum_{i=0}^{n-1} \alpha_{n-1}^{(i)} x_i.$$

Таким чином, отримуємо такі рекурентні співвідношення на $\alpha_n^{(i)}$:

$$\begin{gathered}
n\alpha_n^{(0)} = n\alpha_{n-1}^{(0)}, \\
(n-2)(\alpha_n^{(1)} - \alpha_n^{(0)}) = n\alpha_{n-1}^{(1)}, \alpha_2^{(1)} = 0, \\
(n-4)(\alpha_n^{(2)} - \alpha_n^{(1)} + \alpha_n^{(0)}) = n\alpha_{n-1}^{(2)}, \alpha_4^{(1)} = 0, \\
\dots \\
(n-2i) \sum_{n-2i>0} \alpha_n^{(i)}(-1)^i = n\alpha_{n-1}^{(i)}, \alpha_{2i}^{(i)} = 0.
\end{gathered}$$

Після спрощення отримуємо таку систему рівнянь

$$\begin{cases}
\alpha_n^{(0)} = 1, \\
\alpha_n^{(1)} = n\left(\dfrac{\alpha_{n-1}^{(1)}}{n-2} + \dfrac{\alpha_{n-1}^{(0)}}{n}\right), \\
\alpha_n^{(2)} = n\left(\dfrac{\alpha_{n-1}^{(2)}}{n-4} + \dfrac{\alpha_{n-1}^{(1)}}{n-2}\right), \\
\dots \\
\alpha_n^{(s)} = n\left(\dfrac{\alpha_{n-1}^{(2)}}{n-2s} + \dfrac{\alpha_{n-1}^{(1)}}{n-2(s-1)}\right), \\
\dots
\end{cases}$$

□

Lemma 3.4. *Рекурентне рівняння*

$$x_n = n\left(\frac{x_{n-1}}{n-s} + \frac{g_{n-1}}{n-(s-2)}\right), x(s) = 0,$$

має наступний розв'язок

$$x_n = n^{\underline{s}} \sum_{i=s}^{n-1} \frac{g_i}{i^{\underline{s-1}}\,(i-(s-3))}.$$

Доведення. Маємо

$$\begin{aligned}
n\left(\frac{x_{n-1}}{n-s} + \frac{g_{n-1}}{n-(s-2)}\right) &= n\left(\frac{(n-1)^{\underline{s}}}{n-s}\sum_{i=s}^{n-2}\frac{g_i}{i^{\underline{s-1}}\,(i-(s-3))} + \frac{g_{n-1}}{n-(s-2)}\right) = \\
&= n\left(\frac{(n-1)^{\underline{s}}}{n-s}\sum_{i=s}^{n-2}\frac{g_i}{i^{\underline{s-1}}\,(i-(s-3))} + \frac{(n-1)^{\underline{s-1}}\,g_{n-1}}{((n-1)^{\underline{s-1}})((n-1)-(s-3))}\right) = \\
&= \frac{n(n-1)^{\underline{s}}}{n-s}\sum_{i=s}^{n-2}\frac{g_i}{i^{\underline{s-1}}\,(i-(s-3))} + \frac{n(n-1)^{\underline{s-1}}\,g_{n-1}}{((n-1)^{\underline{s-1}})((n-1)-(s-3))} = \\
&= n^{\underline{s}}\left(\sum_{i=s}^{n-2}\frac{g_i}{i^{\underline{s-1}}\,(i-(s-3))} + \frac{g_{n-1}}{((n-1)^{\underline{s-1}})((n-1)-(s-3))}\right) = \\
&= n^{\underline{s}}\sum_{i=s}^{n-1}\frac{g_i}{i^{\underline{s-1}}\,(i-(s-3))}.
\end{aligned}$$

Тут використали співвідношення

$$\frac{n(n-1)^{\underline{s}}}{n-1} = n^{\underline{s}}, n(n-1)^{\underline{s-1}} = n^{\underline{s}},$$

$\square$

Розв'язуючи рекурентні рівняння при початковій умові $\alpha_{2s}^{(s)} = 0$ послідовно знаходимо

$$\alpha_n^{(0)} = 1,$$

$$\alpha_n^{(1)} = \frac{1}{2}(n-1)(n-2),$$

$$\alpha_n^{(2)} = \frac{1}{6}\,(n-4)\,(n-3)\,(n-2)\,n,$$

$$\alpha_n^{(3)} = \frac{1}{144}\,(n-1)\,n\,(n-4)\,(n-5)\,(7\,n-17)\,(n-6)\,,$$

$$\alpha_n^{(4)} = \frac{1}{2880}\,(n-8)\left(39\,n^2-296\,n+545\right)(n-7)\,(n-6)\,(n-2)\,(n-1)\,n$$

Будемо розглядати невідомі послідовності $\alpha_n^{(s)}, s > 1$ у базисі із спадних факторіалів. Нехай

$$\alpha_n^{(s)} = (n-(2s-1))\left(\beta_0^{(s)} n^{\underline{s-1}} + \beta_1^{(s)} n^{\underline{s}} + \cdots + \beta_s^{(s)} n^{\underline{2s-1}}\right) = (n-(2s-1))\sum_{i=0}^{s} \beta_i^{(s)} n^{\underline{s-1+i}}.$$

Тоді

$$\alpha_n^{(s)} = (n-(2s-1))\sum_{i=0}^{s} \beta_i^{(s)} n^{\underline{s-1+i}} = \sum_{i=0}^{s-1} \beta_i^{(s)} (n-(s+i-1)-(s-i)) n^{\underline{s+i}} =$$

$$= \sum_{i=0}^{s} \beta_i^{(s)} (n-(s+i-1)) n^{\underline{s+i-1}} - \sum_{i=0}^{s} \beta_i^{(s)} (s-i) n^{\underline{s-1+i}} = \sum_{i=0}^{s} \beta_i^{(s)} n^{\underline{s+i}} - \sum_{i=0}^{s-1} (s-i) \beta_i^{(s)} n^{\underline{s-1+i}}$$

З іншого боку ми маємо

$$\alpha_{n-1}^{(s)} = (n-2s)\sum_{i=0}^{s} \beta_i^{(s)} (n-1)^{\underline{s-1+i}},$$

$$\alpha_{n-1}^{(s-1)} = (n-(2s-2))\sum_{i=0}^{s-1} \beta_i^{(s-1)} (n-1)^{\underline{s+i-2}}.$$

Тоді

$$\alpha_n^{(s)} = n\left(\frac{\alpha_{n-1}^{(s)}}{n-2s} + \frac{\alpha_{n-1}^{(s-1)}}{n-(2s-2)}\right) = \sum_{i=0}^{s} n\beta_i^{(s)} (n-1)^{\underline{s-1+i}} + \sum_{i=0}^{s-1} n\beta_i^{(s-1)} (n-1)^{\underline{s-2+i}} =$$

$$= \sum_{i=0}^{s} \beta_i^{(s)} n^{\underline{s+i}} + \sum_{i=0}^{s-1} \beta_i^{(s-1)} n^{\underline{s-1+i}}.$$

Отже

$$\sum_{i=0}^{s-1} \beta_i^{(s-1)} n^{\underline{s-1+i}} = -\sum_{i=0}^{s-1} (s-i) \beta_i^{(s)} n^{\underline{s-1+i}}$$

Звідси, прирівнявши відповідні коефіцієнти, зразу знаходимо, що

$$\beta_i^{(s)}=-\frac{\beta_i^{(s-1)}}{s-i}, i=0\dots s-1.$$

Коефіцієнт $\beta_s^{(s)}$ знайдемо з умови, що $\alpha_{2s}^{(s)}=0$. Маємо

$$\alpha_{2s}^{(s)}=\sum_{i=0}^{s}\beta_i^{(s)}(2s)^{\underline{s-1+i}}=\sum_{i=0}^{s-1}\beta_i^{(s)}(2s)^{\underline{s-1+i}}+\beta_s^{(s)}(2s)^{\underline{2s-1}}=0$$

Звідси, врахувавши $(2s)^{\underline{2s-1}}=2s(2s-1)\dots 2=(2s)!$, отримаємо

$$\beta_s^{(s)}=-\frac{1}{(2s)!}\sum_{i=0}^{s-1}\beta_i^{(s)}(2s)^{\underline{s-1+i}}=-\sum_{i=0}^{s-1}\frac{\beta_i^{(s)}}{(s-i+1)!}.$$

Таким чином, маємо такі рекурентні співвідношення для $\beta_n^{(s)}$:

$$\begin{aligned}
&\beta_0^{(s)}=-\frac{\beta_0^{(s-1)}}{s},\\
&\beta_1^{(s)}=-\frac{\beta_1^{(s-1)}}{s-1},\\
&\beta_2^{(s)}=-\frac{\beta_1^{(s-1)}}{s-2},\\
&\dots\\
&\beta_{s-1}^{(s)}=-\beta_1^{(s-1)},\\
&\beta_s^{(s)}:=b_s=-\sum_{i=0}^{s-1}\frac{\beta_i^{(s)}}{(s-i+1)!}.
\end{aligned}$$

Звідси знаходимо, що

$$\beta_i^{s}=\frac{(-1)^{s-i}}{(s-i)!}b_i,\ \text{for } i=0,\dots,s-1.$$

Отримуємо таке рекурентне співвідношення на послідовність b_s :

$$b_s=-\frac{(-1)^s}{((s+1)!)^2}-\sum_{i=1}^{s-1}\frac{(-1)^{s-i}}{((s-i)(s-i+1)!)}b_i,$$

Поклавши $b_0=1$ отримаємо, що елементи послідовності b_n задовільняють таке рекурентне співвідношення

$$\sum_{i=0}^{s}\frac{(-1)^{n-i}}{(s-i)!(s-i+1)!}b_n=0.$$

Розглянемо ряд

$$\sum_{n=0}^{\infty}\frac{(-1)^n}{n!(n+1)!}z^n=\frac{1}{\sqrt{z}}J_1(\sqrt{4z}\,)$$

і звичайну породжуючу функцію

$$G(b_n,z)=\sum_{n=0}^{\infty}b_nz^n.$$

Тоді, враховуючи рекурентне співвідношення знйдемо, що

$$G(b_n, z)\frac{J_1(\sqrt{4z}\,)}{\sqrt{z}} = 1.$$

Тому

$$G(b_n, z) = \frac{\sqrt{z}}{J_1(\sqrt{4z}\,)} = 1 + \frac{1}{2}\,z + \frac{1}{6}\,z^2 + \frac{7}{144}\,z^3 + \frac{13}{960}\,z^4 + \frac{107}{28800}\,z^5 + \frac{409}{403200}\,z^6 + \cdots +$$

Отже ми довели таку теорему

Theorem 3.2. *A Appel-Fibbonaci ψ_{AF} interwining map has the form*

$$\psi_{AL}(x_n) = x_{n+1} + \alpha_n^{(1)} x_{n-1} + \alpha_n^{(2)} x_{n-3} + \ldots + \alpha_n^{(i)} x_{n+1-2i} + \ldots + \alpha_n^{([\frac{n-1}{2}])} x_{n+1-2[\frac{n-1}{2}]},$$

where

$$\alpha_n^{(s)} = (n-2s+1)\left(\frac{(-1)^s}{s!} b_0 n^{\underline{s-1}} + \cdots + \frac{(-1)^{s-i}}{(s-i)!}\, b_i n^{\underline{s+i}} + \cdots + b_s n^{\underline{2s-1}}\right),$$

and the generating function for $b_0, b_1, \ldots, b_n, \ldots$ are defined через обернену функцію Бесселя

$$\sum_{i=0}^{\infty} b_i z^i = \frac{\sqrt{z}}{J_1(\sqrt{4z}\,)}.$$

Задача. Знайти формулу для b_n.

Із рекурентного співвідношення для многочленів Фібоначчі виплаває такйи зв'язок між диференціюваннями Вейтценбека і Фібоначчі

$$\mathcal{D}(x_n) = 2D_F(x_n) - x_2 D_F(x_{n-1}).$$

Література

[1] Falcon, Sergio; Plaza, Angel On k-Fibonacci sequences and polynomials and their derivatives. Chaos Solitons Fractals 39, No. 3, 1005-1019 (2009)

[2] P. Filipponi and A.F. Horadam. "Derivatives of Fibonacci and Lucas Polynomials" Applications of Fibonacci Numbers. Volume 4. Edited by G.E. Bergum, A.N. Philippou and A.F. Horadam. Dordrecht: Kluwer, 1991: 99-108.

[3] L. Bedratyuk, Semi-invariants of binary forms and identities for Bernoulli, Euler and Hermite polynomials, *Acta Arith.*, 151 (2012), 361-376

[4] Glenn O. Treatise on theory of invariants.–Boston,1915.–312P.

[5] Nowicki A. Polynomial derivation and their Ring of Constants, –Torun:UMK,1994.–170p.

[6] van den Essen A. Polynomial automorphisms and the Jacobian conjecture.–Basel: Birkhäuser. –2000.–329 p.

DEPARTMENT OF MATHEMATICS AND STATISTICS, KHMELNYTSKYI NATIONAL UNIVERSITY, KHMELNYTSKYI, INSTYTUTSKA ,11, 29016, UKRAINE

E-mail address: `leonid.uk@gmail.com`